

Note during this time, the fetus is

Suspended in the uterus is, it does not fall out, but in the direction of raising Chances of pregnancy, relative rest is recommended after three days of resuming the activity normal and everyday life is unhindered be 14 to 17 days after embryo transfer a pregnancy blood test should be done It is better to do this test 2 times with a gap 2 to 3 days after the first test.

If the result of the blood test is positive be: You should take all your medicines according to the previous routine continued for 3 months and 10 days after positive test for ultrasound see your doctor

If the answer to both of your tests if it is negative, you are not pregnant and the kidney Stop taking your medication to get regular Then see your doctor.

At duration of transfer until testing

pay attention to the following points:

Activity: Intense and sudden activity and do not rotate because yet your ovaries are affected by medications Large and bulky ovulation stimulant and there is a possibility of twisting them and too

Embryo transfer

Embryo transfer into the uterus

To transfer the embryo 3 to 5 days after the operation Ovulation with wife at 7 am See the center.

The night before visiting for embryo transfer take a shower.

Before going to the operating room for transfer Avoid going to the bathroom the bladder must be full.

May be the day of embryo transfer, transfer operation for reasons such as not forming an embryo Suitable, not suitable endometrium, accumulation Fluid inside the abdomen, according to the doctor's diagnosis do not treat.

If embryo transfer does not take place and the embryos should be frozen, the next reference should be at least two months later, a few days before the start Menstruation usually

The embryo transferred to your uterus, too it is now suspended in your womb and it takes about three days to get inside the endometrium penetrates the uterus.



University of Medical Sciences
and Health Services, West
Azerbaijan

Kowsar Women's Educational and
Medical Center

Embryo transfer



Contraindicated:

Infertility and stimulation of
ovulation



Thanks for your reading

2024

Sneezing and coughing especially during the transfer operation It is very harmful, so if you catch a cold Corrosion, must take medicine under the supervision of a doctor do it from arbitrarily taking any medicine in during this period, strictly avoid which for some reasons are treated with certain drugs be sure to check the medication list before transferring check your intake with your doctor.

Spotting and bleeding:

In embryo transfer cycles, bleeding and spotting after transfer are common and symptoms

It is not the absence of pregnancy, so in this case, spotting or bleeding, do not stop your medication arbitrarily and consult your doctor or medical staff as soon as possible.



much activity causes irritation Uterine contractions.

Sleep and rest in bed:

Just sleeping in bed after the transfer operation is a misconception and not recommended

Bathing: After the transfer, bathing is allowed.

Sports: from doing sports that need to have a long standing or cause an increase Intra-abdominal pressure or shortness of breath avoid becoming.

Light walking is recommended after the first three days.

Swimming pools and bathtubs are not recommended due to the increased chance of infection.

Try to avoid straining during defecation - do not have intercourse for two weeks.

Attending work and resuming work activities, preferably three days after the transfer without hindrance.

Air and land travel should be as much as possible up to 48 hours after embryo transfer be delayed.